# JEE Journal of Ecological Engineering

Journal of Ecological Engineering, 2025, 26(4), 250–263 https://doi.org/10.12911/22998993/200145 ISSN 2299–8993, License CC-BY 4.0 Received: 2024.12.31 Accepted: 2025.01.31 Published: 2025.02.17

# Nutrient use efficiency in edamame (*Glycine max* L.) production with poultry manure and biofertilizer application

Maya Melati<sup>1\*</sup>, Hayatun Nufuz<sup>2,3</sup>, Alfani Bisri<sup>2</sup>, Dhika Prita Hapsari<sup>1</sup>, Willy Bayuardi Suwarno<sup>1</sup>

- <sup>1</sup> Department of Agronomy and Horticulture, Faculty of Agriculture, IPB University (Bogor Agricultural University), Meranti Street, IPB Darmaga Campus, Bogor 16680, Indonesia
- <sup>2</sup> Study Program of Agronomy and Horticulture, Faculty of Agriculture, IPB University (Bogor Agricultural University), Meranti Street, IPB Darmaga Campus, Bogor 16680, Indonesia
- <sup>3</sup> Agroobot Indonesia
- \* Corresponding author's e-mail: maya\_melati@apps.ipb.ac.id

# ABSTRACT

Nutrient use efficiency (NUE), as one of the agriculture ecological efficiency measures, is an important concept to evaluate the production systems of the crop. NUE in organic farming is less studied. Large quantities of poultry manure are needed in organic farming, therefore biological fertilizer is considered to increase the efficiency of fertilizer use. The research was aimed at investigating the effects of poultry manure rates and biofertilizer on edamame yield and the NUE from two experiments at different times. The experiments were done in 2023 and 2024. Each experiment was arranged in a split plot block design with biofertilizer as the main plot (without and with) and poultry manure rates as subplot (0, 8, 16, and 24 tons ha<sup>-1</sup>), replicated 3 times. A combined analysis of variance from two experiments was done. The study investigated the edamame pod yield, agronomic efficiency based on manure rates and based on N, P, K supplied by manure (AE, AE<sub>N</sub>, AE<sub>P</sub>, AE<sub>K</sub>), and partial factor productivity  $(PFP_{N}, PFP_{\mu}, PFP_{\kappa})$ . The results revealed that filled pod number and weight of aboveground biomass increased with higher manure rates. The effect of interaction between biofertilizer, manure rates, and year was significant on pod weight per plant and edamame pod yield. The increases in pod weight and pod yield following manure rates showed different trends according to the presence or absence of biofertilizer and differences in experimental time. Higher increases in pod weight (141.5–293.3%) and pod yield (125.5–174%), following manure rates, occurred in the first experiment with biofertilizer. In experiment 2, pod yield was not affected by manure rates either with or without biofertilizer. AE, AE<sub>N</sub>, AE<sub>P</sub>, and AE<sub>K</sub> were significantly affected by the interaction of biofertilizer, manure rates, and year of the experiment; they decreased with increasing manure rates in the first experiment. Positive values of AE, AE<sub>N</sub>, AE<sub>p</sub>, and AE<sub>k</sub> were found in the first experiment with biofertilizer. In contrast, in the  $2^{nd}$  experiment, positive values of AE, AE<sub>N</sub>, AE<sub>p</sub> and AE<sub>K</sub> were found without biofertilizer. PFP<sub>N</sub>, PFP<sub>p</sub> and PFP<sub>K</sub> decreased with higher poultry manure rates. Different soil characteristics and methods of biofertilizer application between two experiments may affect the NUE.

Keywords: agronomic efficiency, harvest index, leaf nutrient, organic farming, partial factor productivity.

#### INTRODUCTION

Edamame or vegetable soybean (*Glycine max* L. Merr.) is usually harvested at the R6-R7 stage when pods are still green but the seeds have fully developed. Good nutritional values of edamame that have been identified are isoflavones (Zeipina et al., 2017), dietary fiber (Johnson et al., 1999; Xu et al., 2012), polyunsaturated fatty acids, such

as linoleic acid and linolenic acid (Kumar et al., 2006a). Rich in sugar contents is one of the things that makes edamame popular (Carneiro et al., 2021; Zeipina et al., 2017).

Edamame can be grown with poultry manure as an external source of nutrients. Poultry manure application can be comparable to that of urea in several crops' yields (Lin et al., 2016). The positive influences of poultry manure on Journal of Ecological Engineering 2025, 26(4), 250–263

crop production through improving soil properties, namely increasing soil organic carbon, pH, available P, K, and Zn. The improvement of soil properties was also found in biological characteristics (Kobierski et al., 2017) and physical characteristics such as reduced soil bulk density, temperature, and increased porosity and moisture content (Adeyemo et al., 2019; Agbede et al., 2017; Rasool et al., 2023). Poultry manure as the only external source of nutrients in organic farming has been studied for the production of Sonchus arvensis with 14 tons manure ha-1 (Melati et al., 2021), okra with 18 tons manure ha-1 (Fhonna et al., 2023), and bitter gourd with 30 tons manure ha-1 (Jabary et al., 2023). To obtain the optimum yield of those crops, high dosages of poultry manure had been determined.

Continuous efforts are made to increase fertilizer use efficiency so the doses of fertilizer can be reduced. One way to increase fertilization efficiency is by providing biofertilizers. Biofertilizers are microbial inoculants, derived from plant roots and root zones, that can be applied to seeds, plant surfaces, or soil. The roles of biofertilizers in plant production are to improve soil fertility, increase nutrient availability, improve plant growth, and increase plant productivity. Another important role of biofertilizer is its ability to protect plants from pests and diseases and ameliorate biotic and abiotic stresses (Kumar et al., 2022; Nosheen et al., 2021). Biofertilizers can increase the availability of N, P, and K (Mitter et al., 2021; Ortega Pérez et al., 2023). The role of biofertilizers in plant growth promoting can contribute to the development of root hairs which in turn can improve water uptake (Magana et al., 2020; Singh et al., 2020; Weber et al., 2018).

Nutrient use efficiency (NUE) or fertilizer use efficiency is an important concept to evaluate the production systems of the crop (Gajanand et al., 2020). Several indices can be used to measure the nutrient use efficiency from the applied fertilizer to assess the response of a crop to a nutrient, they are recovery efficiency (RE), physiological efficiency (PE), internal utilization (IE), agronomic efficiency (AE), and partial factor productivity (PFP) (Dobermann, 2007). Various nutrient use efficiency (NUE) indices can also be grouped into fertilizer-based, plant-based, soil-based, isotope-based, ecology-based, and system-based. Each of the indexes has strengths and limitations to use in determining the nutrient use efficiency of the crops (Congreves et al., 2021). A similar calculation of nitrogen use efficiency (NUE) can be used to determine different nutrients, for example, phosphorus use efficiency (PUE) and other nutrients. The nutrient use efficiency is not only for calculating the efficiency due to fertilizer application (Mandic et al., 2015; Schütz et al., 2018) but can also as a result of crop cultivation arrangement (intercropping) (Suntari et al., 2023; Zhang et al., 2015). Improvement of NUE is also aimed at reducing the potential for excess fertilizer to pollute the environment.

Because of its nutritional value and potential health benefits, the consumption of edamame has increased not only in Asia but also in the US (Yu et al., 2021). To meet the needs for edamame, the production needs to be increased, including through organic cultivation by using organic fertilizer. It is important to pay attention to the efficiency of inputs both for economic reason and concern for the environment. In organic farming practices, that apply large amounts of organic fertilizer, the effects of biofertilizer on nutrient efficiency have not been revealed yet. Therefore, the current study was aimed at investigating the effects of four levels of poultry manure and the application of biofertilizer on edamame soybean yield traits and the nutrient use efficiency from two experiments at different times. Nutrient use efficiencies were determined based on agronomic efficiency (AE) and partial factor productivity (PFP).

# MATERIALS AND METHODS

Two experiments at different times have been conducted. The 1<sup>st</sup> experiment was carried out from January to April 2023, while the 2<sup>nd</sup> experiment was in November 2023-March 2024 at IPB Experimental station in Cikarawang, Darmaga, Bogor, Indonesia (-6.5497175° S, 106.7287026° E). The region has a tropical climate with annual precipitation of 2000–3000 mm. The experimental site had tropical latosol soil (Alfisol). The chemical properties of the soil were analyzed before the application of manure (Table 1).

The experiment used edamame soybean seed from Biomax 1 variety, dolomite, laying hen manure, and commercial biological fertilizer (consortium of *Bacillus polymyxa, Pseudomonas flurescens, Rhizobium* sp). The chemical traits of poultry manure in presented in Table 2.

Soil properties	2023	2024
pH H <sub>2</sub> O	6.19	6.28
C-organic (%)	1.86	1.66
N-total (%)	0.25	0.22
P-available (ppm P205)	30.87	49.54
CEC (cmol/kg)	17.42	15.71
K-exc (cmol K/kg)	0.22	0.28
P-potential (mg P2O5/100g)	107.26	174.54
K-potential (mg K <sub>2</sub> O/100 <sub>g</sub> )	29.77	36.09

Table 1. Chemical properties of soil before manure application

 Table 2. Chemical properties of poultry manure applied in 2023 and 2024 and potential supply of nutrients for each dose of manure application

		Potenti	al supply o from man	of nutrients ure		Potential supply of nutrient from manure			
Manure properties	2023	8 tons	16 tons	6 tons 24 tons 20		8 tons	16 tons	24 tons	
			(kg∙ha⁻¹	)		(kg·ha-1)			
рН	7.31				8.09				
Moisture content	22.53 %				18.76 %				
C-organic	17.48 %				23.14 %				
N-total	1.43 %	88.6	177.3	265.9	2.14 %	139.1	278.2	417.2	
P2O5 total	5.26 %	325.9	651.9	977.9	3.84 %	249.6	499.1	748.7	
K <sub>2</sub> O total	3.70 %	229.3	458.6	687.9	3.41 %	266.5	532.9	799.4	

# **Experimental design**

Experiments 1 and 2 applied a split-plot design with biological fertilizer (bio-fertilizer) as the main plot (with and without biofertilizer) and manure rates as sub-plot (0, 8, 16, and 24 tons  $\cdot$  ha<sup>-1</sup>). Each treatment was replicated 3 times. The difference between experiment 1 and 2 was the method of biofertilizer application.

# Land preparation

Following the land preparation, dolomite lime and laying hen manure were spread over the soil surface and then mixed. Each plot size was  $3 \times 1.25$  m (experiment 1) and  $3 \times 1.5$  m (experiment 2). Each plot was separated by a distance of 50 cm.

# Seed planting, application of bio-fertilizer, and harvest

Soybean seeds were planted with a  $25 \times 20$  cm distance, 2 seeds per planting hole. In experiment 1, seeds were soaked for 15 minutes in a biofer-tilizer solution with a concentration of 10 g·L<sup>-1</sup>

water, following the instructions of the product. In experiment 2, seeds were soaked in a biofertilizer solution with a concentration of 10 g·L<sup>-1</sup> water before planting. Additional application of biofertilizer was done twice. The first application was after planting, then at the V3 plant growth phase, with the concentration of 5 g·L<sup>-1</sup> water as much as 100 mL per plant. Soybean pods were harvested immature between the R6 and R7 reproductive stages. At this stage, the pods are still green but the seeds have fully developed. At harvest, soybean plants were uprooted and then separated into pods and biomass.

# Sample collection, measurements, and calculations

Ten plants were sampled to determine the pod number per plant, pod weight per plant, and fresh weight of plant biomass at harvest 70–72 days after planting (R6 stage). The harvest index represents the ratio between the plant product and biomass (Eq. 1). To measure the nutrient efficiency, agronomic efficiency was calculated with Eq. 2–5, while the partial factor productivity was determined by Eq. 6–8 (Dobermann, 2007).

$$HI = \frac{WFPP(g)}{FWABH + PW(g)}$$
(1)

where: *HI* – harvest index, *WFPP* – weight of filled pod per plant, *FWABH* – fresh weight of aboveground biomass at harvest, *PW* – pod weight.

$$AE = \frac{Ym - Yo(ton)}{Manure applied(ton)}$$
(2)

where: Ym – yield with manure, Yo – yield without manure.

$$(AE_N) = \frac{Ym - Yo(kg)}{N \text{ supplied by manure } (kg)}$$
(3)

where:  $AE_N - N$  - based agronomic efficiency, Ym - yield with manure, Yo - yield without manure.

$$(AE_P) = \frac{Ym - Yo(kg)}{P \text{ supplied by manure } (kg)}$$
(4)

where: AEp - P – based agronomic efficiency, Ym – yield with manure, Yo – yield without manure.

$$(AE_K) = \frac{Ym - Yo(kg)}{K \text{ supplied by manure } (kg)}$$
(5)

where:  $AE_{K} - K$  - based agronomic efficiency, Ym - yield with manure, Yo - yield without manure.

$$(PFP_N) = \frac{Pod \ yield \ (kg \cdot ha^{-1})}{N \ supplied \ from \ manure \ (kg \cdot ha^{-1})} \quad (6)$$

where:  $(PFP_N)$ -N-based partial factor productivity.

$$(PFP_P) = \frac{Pod yield (kg \cdot ha^{-1})}{P \text{ supplied from manure } (kg \cdot ha^{-1})} \quad (7)$$

where: (*PFPp*) – *P* – based partial factor productivity.

$$(PFP_K) = \frac{Pod yield (kg \cdot ha^{-1})}{K \text{ supplied from manure } (kg \cdot ha^{-1})} \quad (8)$$

where:  $(PFP_{K})-K$ -based partial factor productivity.

The concentrations of N, P, and K in leaf tissues were determined from samples collected at the V3 growth stage. Leaves from two plants of each plot were oven-dried at 60 °C. Leaf N was determined with Kjeldahl, P with spectrophotometer UV-VIS, and K with atomic adsorption spectrophotometer (AAS). Analysis of N, P, and K in leaves, was done on treatment with biofertilizer.

#### **Statistical analysis**

The data were processed with analysis of variance using the statistical tool for agricultural

research (STAR). A combined analysis of variance from two experiments was done with a Split Plot in Randomized Block Design. Three factors were used as sources of variance namely year (2023 and 2024), biofertilizer (without and with), and manure rates (0, 8, 16, 24 tons  $\cdot$  ha<sup>-1</sup>). The statistically significant differences between means were determined by the LSD test and the significance level was set at p  $\leq 0.05$ .

Data with concentrations of N, P, and K in leaves were analyzed by combined analysis of variance from two experiments in Randomized Block Design. Two factors were used as sources of variance namely year (2023 and 2024) and manure rates (0, 8, 16, 24 tons ha<sup>-1</sup>).

# RESULTS

#### Plant biomass

Based on the result of the combined analysis of variance from the two experiments, attention was given to sources of variability that refer to the pooled error with a degree of freedom that is more than 6. The effects of manure rates were significant at p < 0.01 on the variables of plant biomass at 5 weeks after planting (5 WAP) and at harvest (9 WAP). Compared to control, the application of poultry manure increased pooled means of fresh and dry weight of aboveground biomass by 43.5-94.9% but there was no significant difference among manure rates. Poultry manure as much as 8, 16, and 24 tons ha<sup>-1</sup> had increased the fresh weight of aboveground biomass at harvest by 24.4, 63.1, and 80.6% compared to control. The biomass weights were different among manure rates (Table 3).

#### **Production components**

For yield attributes, the significant effects of manure rates were found on filled pod number (p<0.01) and harvest index (p<0.05) (Table 4). The application of poultry manure with the rate of 16 and 24 tons  $\cdot$  ha<sup>-1</sup> increased the filled pod number by 64.2% compared to the control (without manure). The highest manure rate resulted in a decrease in harvest index by 6.8% compared to control (Table 4).

The significant effect of the interaction of year-biofertilizer-manure rates was found in pod weight per plant (p<0.01) and pod yield (p<0.05)

Treatments	Fresh wei	ght of abov 5 WA	reground bior P (g)	nass at	Dry weight of aboveground biomass at 5 WAP (g)					Fresh weight of aboveground biomass at harvest 9 WAP (g)			
	2023	2024	pooled		2023	2024	pooled		2023	2024	pooled		
Biofertilizer													
+ biofertilizer	34.61	27.64	31.02		9.05	5.61	7.23		18.58	59.10	38.84		
- biofertilizer	38.23	26.99	32.37		9.46	5.67	7.45		18.42	50.00	34.21		
Manure rates (ton·ha <sup>-1</sup> )													
0	20.72	20.42	20.57	b	5.18	4.30	4.73	b	9.23	42.22	25.72	d	
8	31.69	27.45	29.53	ab	8.40	6.11	7.21	а	13.62	50.37	31.99	с	
16	51.21	28.98	39.32	а	13.23	5.91	9.23	а	24.21	59.67	41.94	b	
24	46.10	33.18	39.37	а	11.19	6.34	8.61	а	26.94	65.94	46.44	а	

**Table 3.** The means of fresh weight and dry weight of aboveground biomass with different manure rates and biofertilizer treatments from the two experiments

Note: Numbers in the same column followed by different letters indicate significant differences based on LSD test at  $p \le 0.05$ .

**Table 4.** The means of filled pod number and harvest index with different manure rates and biofertilizer treatments

 from the two experiments

Transferration	F	illed pod numbe	er per plant	Harvest index				
Treatments	2023	2024	pooled		2023	2024	pooled	
Biofertilizer								
+ biofertilizer	18.7	24.0	21.4		0.66	0.52	0.59	
- biofertilizer	17.4	20.9	19.2		0.68	0.54	0.61	
Manure rates (ton · ha · 1)								
0	11.4	18.2	14.8	b	0.68	0.55	0.61	а
8	14.7	20.8	17.8	b	0.71	0.54	0.62	а
16	22.1	26.4	24.3	а	0.66	0.54	0.60	ab
24	23.9	24.7	24.3	а	0.64	0.50	0.57	b

Note: Numbers followed by different letters in the same column indicate significant differences based on LSD test at  $p \le 0.05$ .

(Fig. 1a and 1b). There are different trends of pod weight in response to the treatments. Without poultry manure but added with biofertilizer, pod weight per plant was 15.15 g (2023) and 56.36 g (2024). When poultry manure was added with biofertilizer, pod weight per plant from experiment 1 (2023) increased between 141.5–293.3%, while from experiment 2 (2024) pod weight increased between 16.4–25.4% as a result of manure application 8–24 tons ha<sup>-1</sup>, compared to without manure (Fig. 1a).

In contrast, without biofertilizer and poultry manure, pod weight per plant was 28.31 g (2023) and 43.82 g (2024). In the application of poultry manure without the addition of biofertilizer, the pod weight of edamame increased between

4.59–87.6% (2023) and 15.5–52.83% (2024) compared to the control without manure (Fig. 1a).

The interaction of year-biofertilizer-manure rates affects the pod yield of edamame. Different trends of pod yield in response to the treatments were found. Without manure but applied with biofertilizer, pod yield was 4.27 tons ha<sup>-1</sup> (experiment 1) and 10.69 tons ha<sup>-1</sup> (experiment 2). With the addition of poultry manure 8–24 tons ha<sup>-1</sup>, edamame pod yield increased between 125.5–174% in experiment 1, but decreased by 1.03–8.04% in experiment 2, compared to without manure (Fig. 1b). However, pod yields were not different among manure rates in experiment 2. Without biofertilizer and poultry manure, pod yield was 7.65 tons ha<sup>-1</sup> (2023) and 9.49 tons ha<sup>-1</sup> (2024). The application of poultry manure 8–24



Figure 1. Interaction effects of manure rates, biofertilizer, and time of experiment on means of pod weight per plant (a) and pod yield (b). Different letters indicate significant differences based on LSD test at  $p \le 0.05$ 

tons ha<sup>-1</sup> increased pod yield by 4.97–44.97% (2023) and 15.6–22.13% (2024) compared to control without manure (Fig. 1b). However, pod yields were not different among manure rates in Experiment 2024.

#### Agronomic efficiency

The effect of the interaction of year-biofertilizer-manure rates was significant on agronomic efficiency (AE) (p<0.01), AE based on N supplied by manure (AE<sub>N</sub>) (p<0.01), AE based on P supplied by manure  $(AE_{p})$  (p<0.01), and AE based on K supplied by manure (AE<sub>k</sub>) (p<0.05). In experiment 1, with biofertilizer, the application of 8 tons of manure ha-1 resulted in agronomic efficiency of as much as 0.623 tons of pod yield/ton manure (Fig. 2a). Increasing manure rates had increased pod yield (Fig. 1b) but the AE became lower. Given the 24 tons of manure ha-1, the AE decreased by 94% compared to AE from 8 tons ha<sup>-1</sup> (Fig. 2a). In experiment 2, with biofertilizer, the application of 8 tons manure ha-1 caused an agronomic efficiency of as much as -0.085 tons pod yield/ton manure (Fig. 2a) or there was a decrease in AE due to the slight reduction of yield (Fig. 1b). The reduction of AE became smaller with increasing manure rates. The AE was not different among 8, 16, and 24 tons of manure  $\cdot$  ha<sup>-1</sup>.

Without biofertilizer, the AE was 0.047 (2023) and 0.238 (2024) tons pod yield/ton manure. In 2023, the AE was not different among 8, 16, and 24 tons of manure  $ha^{-1}$ , but in 2024, the AE with 24 tons of manure decreased by 41% compared to AE from 8 tons  $ha^{-1}$  (Fig. 2a). Based on the amount of N, P, and K potentially

supplied by poultry manure, the agronomic efficiency was determined for each nutrient. In experiment 1, positive values of  $AE_N$  (0.060 kg pod yield/kg N supplied),  $AE_P$  (0.017 kg·pod yield/kg·P supplied), and  $AE_K$  (0.023 kg·pod yield/kg·K supplied) were found from the application of 8 tons·manure·ha<sup>-1</sup> with biofertilizer. Lower  $AE_N$ ,  $AE_P$ , and  $AE_K$  were found with increasing manure rates (Fig. 2b, 2c, 2d). The values of  $AE_N$ ,  $AE_P$ , and  $AE_K$  decreased by 9.2, 2.57, and 4.32% respectively.

Still in experiment 1, without biofertilizer, positive values of AE<sub>N</sub> (0.004 kg·pod yield/kg·N supplied),  $AE_{P}(0.001 \text{ kg} \cdot \text{pod yield/kg} \cdot \text{P supplied})$ , and  $AE_{\nu}$  (0.002 kg·pod yield/kg K supplied) were found from the application of 8 tons manure ha-1 (Fig. 2b, 2c, 2d), but they were smaller than those with biofertilizer. The  $AE_N$ ,  $AE_P$ , and  $AE_K$  without biofertilizer were not different among 8, 16, and 24 tons of manure ha-1. Different trends were shown by experiment 2. Negative values of  $AE_{y}$ (-0.003 kg pod yield/kg·N supplied),  $AE_{p}$  (-0.002 kg·pod yield/kg·P supplied), and  $AE_{\kappa}$  (-0.002 kg·pod yield/kg·K supplied) were found from the application of 8 ton manure ha-1 with biofertilizer. On the contrary, without biofertilizer, positive values were found for AE<sub>N</sub> (0.010 kg·pod yield/ kg·N supplied), AE<sub>p</sub> (0.007 kg pod yield/kg·P supplied), and  $AE_{K}$  (0.007 kg·pod yield/kg·K supplied). These  $AE_{N}$ ,  $AE_{p}$  and  $AE_{K}$  were not different among manure rates.

# Partial factor productivity

Manure rates significantly affect the partial factor productivity of N (PFP<sub>N</sub>) (p<0.001)



Figure 2. Interaction effects of manure rates, biofertilizer, and time of experiment on agronomic efficiency (AE) based on manure dose (a), AE based on N- (b), AE based on P- (c), and AE based on K- (d) supplied from manure. Different letters indicate significantly different based on LSD test at p ≤ 0.05

and of K (PFP<sub>K)</sub> (p<0.01), while P (PFP<sub>P</sub>) was influenced by an interaction between year and manure rates (Table 5). PFP<sub>N</sub>, PFP<sub>P</sub>, and PFP<sub>K</sub> decreased with increasing manure rates. Given 24 tons of manure ha<sup>-1</sup>, PFP<sub>N</sub>, PFP<sub>P</sub>, and PFP<sub>K</sub> decreased by 61.2, 57, 66.6, and 61.9%, respectively, compared to 8 tons of manure ha<sup>-1</sup>. PFP<sub>P</sub> from experiment 2 was higher than in 2023.

#### Leaf nutrient content

Combined analysis of variance from two experiments for data of concentration of N, P, and K in leaves resulted in significant effects of manure rate on leaf K (p<0.05) (Table 5). Nitrogen and phosphorus contents in soybean leaves were not different among manure rates, while the potassium content

						_							
Tractmonto	PFP-N				PFP-P					PFP-K			
Treatments	2023	2024	Pooled		2023		2024		Pooled	2023	2024	Pooled	
Biofertilizer													
+ biofertilizer	72.23	44.49	58.36		19.64		24.79		22.22	27.92	23.22	25.57	
Biofertilizer	64.59	49.96	57.28		17.56		27.84		22.70	24.96	26.08	25.52	
Manure rates (tonne·ha <sup>.1</sup> )													
8	99.6	77.3	88.5	а	27.1	а	43.1	а	35.1	38.5	40.4	39.4	a
16	62.7	38.5	50.6	b	17.1	b	21.5	b	19.3	24.3	20.1	22.2	b
24	42.9	25.8	34.3	с	11.7	с	14.4	с	13.0	16.6	13.5	15.0	с

Table 5. Effects of manure rates on partial factor productivity (PFP) based on N, P, K applied from manure

Note: Numbers in the same column followed by different letters indicate significant differences based on LSD test at  $p \le 0.05$ .

was significantly affected by manure rates (Table 6). The application of 16 tons of manure  $ha^{-1}$  increases K by 42% compared to control.

#### **Correlation among variables**

The correlation matrix revealed that pod yield positively correlated with weight of aboveground biomass at 5 weeks after planting (V3 growth phase) (p<0.001). This correlation shows that the plant performance at V3 growth phase can determine the edamame pod production. This correlation matrix also shows that the agronomic efficiency AE, AE-N, and AE-P, positively correlated to pod yield which means that the higher the agronomic efficiency, the higher the yield of edamame (Fig. 3).

#### DISCUSSION

#### The impact of poultry manure application

The current study showed that the use of poultry manure as a source of nutrients in organic farming can support the production of edamame soybeans. The increase in manure rates increases the potential supply of nutrients to the soybean plants (Table 2) and results in the increase of N and P in leaf tissue is not significant, and K in leaf tissues (significantly increases by 42%) (Table 6). Nitrogen, phosphorus, and potassium are the three major nutrients required by plants. Nitrogen is the main essential nutrient for plants, a component of the protein. The important role of nitrogen



**Figure 3.** Correlation matrix depicting relationships between edamame pod yield and different variables FW\_5: fresh weight of aboveground biomass at 5 weeks after planting; DW\_5: dry weight of aboveground biomass at 5 weeks after planting, FW\_9: fresh weight of aboveground biomass at harvest (9 WAP); PN: filled pod number; PW: pod weight per plant; PY: pod yield; HI: harvest index; AE: agronomic efficiency based on manure; AE\_N: AE based on N supplied from manure; AE\_P: AE based on P supplied from manure; AE\_K: AE based on K supplied from manure; PFP\_N: nitrogen use efficiency; PFP\_P: phosphorus use efficiency; PFP\_K: potassium use efficiency

Manure rates		Leaf N (%	)		Leaf P (%	6)	Leaf K (%)			
(ton · ha⁻¹)	2023	2024	Pooled	2023	2024	Pooled	2023	2024	Pooled	
0	4.30	4.04	4.17	0.32	0.38	0.35	0.90	1.73	1.31	b
8	4.35	3.85	4.10	0.33	0.37	0.35	1.37	1.72	1.54	ab
16	4.43	4.15	4.29	0.32	0.38	0.35	1.74	1.98	1.86	а
24	4.65	4.29	4.47	0.32	0.40	0.36	1.73	1.94	1.84	а

Table 6. Means of concentration of N, P, and K in soybean leaves with different manure rates

**Notes:** Numbers followed by different letters in the same column indicate significant differences based on LSD test at  $p \le 0.05$ .

in plant physiology is because N is the major component of chlorophyll, so it determines the photosynthetic rate of the plant which in turn the plant productivity (Fathi, 2022). In the process of plant growth, nitrogen controls cell division and cell elongation (Luo et al., 2020; Roggatz et al., 1999). Phosphorus is a structural component of nucleic acid, as a component of ADP, and functions in carbohydrate transfer (Hawkesford et al., 2012). Unlike nitrogen and phosphorus, potassium is a free cation and not a component of molecular structure, but K also plays a key role in plant growth and photosynthetic (Sardans & Peñuelas, 2021), especially as osmoregulation which is important for stomata movement and cell extension (Hawkesford et al., 2012). In the current experiment, the supply of N, P, and K may have improved the photosynthetic of the plants, therefore increasing the aboveground biomass at 5 weeks after planting (Table 3). The aboveground biomass at Vn stage positively correlated to the pod yield and indicates that the increase in plant biomass has increased pod yield.

Poultry manure contains N, P, and K and the potential supply of nutrients (Table 2) from each manure dose may have improved the soil properties. The organic carbon from poultry manure, which is in good values (18-22%) (Table 2) can increase the soil organic carbon. The soil organic carbon accumulation can accelerate the formation of soil aggregation (Zhao et al., 2023). Improvements in soil nutrient availability as a result of organic fertilizer application, are caused by improvements in the biological, physical, and chemical properties of the soil (De Sousa Lima et al., 2021). Poultry manure affects soil properties by increasing soil pH, soil organic carbon, and available P, exchangeable Ca, Mg, and K. It has been reported earlier that increasing poultry manure doses increased the concentration of N, P, and K in leaf tissues of soybean (Soremi et al., 2017), and increased grain yield of soybean (Ahmadi & Arien, 2022). The application of poultry manure can be considered environmentally friendly because the organic fertilizers release nutrients slowly (Asadu et al., 2024).

#### **Edamame productivity**

The interaction effect of year-biofertilizermanure rates affects filled pod weight per plant and pod yield (Fig. 1). Different times of experiment showed different responses of pod weight and pod yield to manure rate treatments with or without biofertilizer. In experiment 1, the increase of pod weight per plant and pod yield were higher when combined with biofertilizer than without biofertilizer (Fig. 1). On the contrary in 2024, the higher increase of pod weight and pod yield with the absence of biofertilizer. Figure 1 shows that all values of pod weight are higher in 2024 than in 2023. The results may relate to the soil properties of these two experiments (Table 1). There are some improvements in soil characteristics in 2024, namely slightly higher soil pH, higher total P and available P, and higher total K and exchangeable K. Better soil characteristics before the initial experiment in 2024 may have contributed to the better response of the plants to the application of poultry manure which resulted in higher biomass then pod weight per plant in 2024 than 2023. It was reported earlier that the influence of residue from previous manure increased wheat productivity (Dhaliwal et al., 2023), while residual compost increased radish yield especially in the area with high soil fertility (Lanna et al., 2018).

In experiment1, pod weight per plant and pod yield increased significantly with the addition of poultry manure combined with biofertilizer. The effect of biofertilizer, which is a consortium of *Bacillus polymyxa* (*Paenibacillus polymyxa*), *Pseudomonas flurescens*, *Rhizobium* sp, may have contributed to promoting plant growth due to the increase of soil nutrient availability and phytohormone. Paenibacillus polymyxa, is a gram-positive bacterium, that has a broad host range so it is a potential PGPR that can promote plant growth (Jeong et al., 2019; Timmusk et al., 2005). Pseudomonas fluorescens is a gram-negative bacteria and one of the genera that are capable of solubilizing insoluble soil P (Anand et al., 2016) into available form (Mitter et al., 2021) and it can promote plant growth (Sahu et al., 2018). Rhizobium is a symbiotic N2-fixing organism that can convert N<sub>2</sub> from the atmosphere into NH, (Yang et al., 2022), promote nodulation, nitrogen fixation, nutrient uptake, plant growth, and seed yield of soybean (Htwe et al., 2019). The positive effect of the combination of poultry manure and biofertilizer has been reported to increase the yield of Stevia rebaudiana (Lozano-Contreras et al., 2021).

The effect of biofertilizers on edamame production may be more prominent in the experiment in 2023 than in 2024. The current study showed that the soil properties before the initial experiment in 2023 had slightly lower soil pH, lower total P, available P, total K, and exchangeable K than in 2024. This soil status may cause plants to get better benefits from the presence of biofertilizers. Several factors determine the efficacy of biofertilizers that can be grouped into edaphic and environmental, inoculant-related, and plantrelated factors (Malusà et al., 2016). In edaphic/ environmental factors, usually under low nutrient availability, plant growth and yield respond better to biofertilization (Da Costa et al., 2014) indicating that the plant gets the maximum benefit from the added microbes (Mitter et al., 2021). Previous studies reported that high soil N inhibits the nodulation of rhizobia (Thilakarathna & Raizada, 2017), while high soil P limits the success of AMF colonization (Jansa et al., 2009).

The different trends of plant's response to the manure rates in 2023 and 2024 can relate to several possibilities. First, there is a residual effect of poultry manure from experiment 2023. Secondly, different methods of microbe inoculation may determine the efficacy of biofertilizers. From inoculant-related factors, there are several critical points for the success of inoculation. The types of microbe, the concentration and formulation of biofertilizer (Mitter et al., 2021). The application method of biofertilizer (on seed or into the soil) and the time and frequency of application (Parnell et al., 2016).

In the current study in experiment 1, biofertilizer was inoculated on the seed, while in experiment 2024, biofertilizer was delivered on the seed and into the soil (twice). The application of biofertilizer as seed coating may have resulted in better root colonizing and facilitated better nutrient supplies to the plants (Mujeeb et al., 2022), but the combination of application methods can also improve plant growth (Abdiani et al., 2019; Mujeeb et al., 2022). However, it is not yet known whether the application of biofertilizers 3 times in experiment 2024 has an impact on the indigenous microorganisms. The impact of biofertilizers on soil microbial diversity is generally positive (Samantaray et al., 2024) but the possible antagonistic interaction is also being concern (Mitter et al., 2021).

# Nutrient use efficiency

In the present study, nutrient use efficiency is presented as agronomic efficiency based on manure rates (AE) and based on the amount of N (AE<sub>N</sub>), P (AE<sub>P</sub>), and K (AE<sub>K</sub>) potentially supplied by the added poultry manure (Fig. 3). Different times of experiment show different trends of AE, AE<sub>N</sub>, AE<sub>P</sub>, and AE<sub>K</sub>.

The nutrient use efficiency, measured as agronomic efficiency (AE, AE<sub>N</sub>, AE<sub>P</sub>, and AE<sub>K</sub>), was found higher in experiment 1 where biofertilizer was combined with the application of poultry manure, compared without biofertilizer. Without biofertilizer, the with 16 and 24 tons of manure were not significantly different over 8 tons (Fig. 3). This shows that there is a positive effect of a combination of poultry manure and biofertilizer on the productivity of edamame soybeans. Generally, organic fertilizers have a positive effect on rhizosphere microorganisms (Malusà et al., 2016). The consortium of Bacillus polymyxa (Paenibacillus polymyxa), Pseudomonas flurescens, Rhizobium sp, may have contributed to promoting plant growth due to the increase of soil nutrient availability and phytohormone.

In experiment 1, combined with the application of biofertilizer, the AE decreased significantly at 24 tons of manure ha<sup>-1</sup>, while the AE<sub>N</sub>, AE<sub>P</sub>, and AE<sub>K</sub> decreased significantly at 16 ton of manure ha<sup>-1</sup> over 8 tons ha<sup>-1</sup>. The effect of biofertilizer, which is a consortium of *Bacillus polymyxa* (*Paenibacillus polymyxa*), *Pseudomonas flurescens, Rhizobium* sp, may have contributed to the increase of soil nutrient availability. Therefore, a further increase in manure rates does not increase production as much as the difference between low rate of manure and no fertilizer. Several factors can cause low nutrient use efficiency, they are leaching loss, gaseous losses, immobilization of fertilizer nutrients, the interaction between different fertilizers, physical and chemical characteristics of soil, soil temperature that influence uptake of N, P, and K, soil moisture, and the fertilizer characteristic itself (Mishra et al., 2023). In line with the current study, it has been previously reported that higher fertilizer rates resulted in lower nutrient (N) use efficiency in common beans (Argaw et al., 2015), and wheat (Sharma et al., 2022). In the current study, the nutrient use efficiency values decreased with higher manure rates because they measured the increase in edamame production between fertilizer and control. However, it can be seen that edamame yield increases more with biofertilizers (Fig. 1) which indicates biofertilizers increase the efficiency of manure application. The positive role of biofertilizer has also been reported previously that the application of plant growth-promoting bacteria (PGPB) can reduce the poultry manure dose from 18 to 12 tons ha-1 for okra fruit production (Magana, 2020).

Several approaches can be considered to improve nutrient use efficiency in organic farming including to improve plant morphological dan physiological characteristics and cultivation techniques. The agronomical approach is essential to improve the nutrient use efficiency in plants to make them economically more feasible as well as to prevent environmental damage due to excess fertilizer. Improvement in cultivation technique including the use of different types of fertilizer (e.g. bio-fertilizers, granule organic fertilizer), nutrient management improvement, modern technology (e.g. precision farming by determining the fertilizer recommendation based on soil testing), and agronomic practices (e.g. cropping pattern) (He et al., 2009; Javed et al., 2022; Mishra et al., 2023).

Different trends of AE,  $AE_N$ ,  $AE_P$ , and  $AE_K$ were shown in experiment 2. Only AE significantly decreased with higher manure rates. In contrast to experiment 1, the combination of poultry manure and biofertilizer did not have a positive effect on AE,  $AE_N$ ,  $AE_P$ , and  $AE_K$  of experiment 2. The efficacy of biofertilizers is determined by, among others, the plant physiological traits and soil characteristics (Malusà et al., 2016). Plants can modify the compound released from the roots and select the specific bacterial communities (Marschner et al., 2004). Root exudates are important to initiate the effect of the rhizosphere in seedling and lateral root emergence. Therefore, to increase the efficacy of biofertilizer, biofertilizer should be delivered on seeds and seedlings (Malusà et al., 2016). In experiment 1, biofertilizer was applied to seeds, while in experiment 2, biofertilizer was delivered to the seed and into the soil (twice). The difference in delivering biofertilizer may have impacted the different efficacy of biofertilizer when combined with poultry manure in 2024 although repeated applications of biofertilizer during the growing season of the plant have been recommended (Malusà et al., 2016).

Partial factor productivity  $PFP_N$ ,  $PFP_P$ , and  $PFP_K$  decreased with higher manure rates (Table 7). The PFP can be used to provide an idea of how production can be increased or maintained with minimal input of nutrients (Congreves et al., 2021). The present study showed that to increase edamame productivity, the nutrient supply from poultry manure cannot be minimized yet.

# CONCLUSIONS

The study revealed that poultry manure, as a source of nutrients, can support organic edamame soybean production in the two experiments. A combination of poultry manure and biofertilizer determines the different responses of soybean plants from different times of experiment. Higher rates of manure increased pod yield with biofertilizer than without biofertilizer in the first experiment but not in the subsequent experiment. Agronomic efficiency was higher with biofertilizer than without biofertilizer in the 1st experiment but vice versa in the 2<sup>nd</sup> experiment. In general, AE decreased with higher manure rates. Partial factor productivity decreased with increasing rates of poultry manure. It is important to increase nutrient use efficiency in organic farming including by looking for more suitable types of microbes or by combining various nutrient sources. The study shows that the application of poultry manure in organic farming needs to be combined with biofertilizer, especially on land that has not accumulated high levels of organic matter. However, in the next planting season, the application method of biofertilizer, in combination with poultry manure, needs to be carefully reviewed. The current study showed that 16 tons ha-1 of poultry manure is sufficient for edamame production with the addition of biofertilizer.

# REFERENCES

- Abdiani, S., Kakar, K., Gulab, G., & Aryan, S. (2019). Influence of biofertilizer application methods on growth and yield performances of green pepper. *International Journal of Innovative Research and Scientific Studies*, 2(4), 45–49. https://doi. org/10.53894/ijirss.v2i4.23
- Adeyemo, A. J., Akingbola, O. O., & Ojeniyi, S. O. (2019). Effects of poultry manure on soil infiltration, organic matter contents and maize performance on two contrasting degraded alfisols in southwestern Nigeria. *International Journal of Recycling of Organic Waste in Agriculture*, 8(Suppl1), S73–S80. https://doi.org/10.1007/s40093-019-0273-7
- Agbede, T. M., Adekiya, A. O., & Eifediyi, E. K. (2017). Impact of poultry manure and NPK fertilizer on soil physical properties and growth and yield of carrot. *Journal of Horticultural Research*, 25(1), 81–88. https://doi.org/10.1515/johr-2017-0009
- Ahmadi, A. Y., & Arien, M. (2022). Poultry manure effects on yield and some agronomic components of Soybean (Glycine max L.) under Khost agro-ecological conditions, Afghanistan. *International Journal of Agriculture Environment and Food Sciences*, 6(1), 1–6. https://doi.org/10.31015/jaefs.2022.1.1
- Anand, K., Kumari, B., & Mallick, M. A. (2016). Phosphate solubilizing microbes: An effective and alternative approach as biofertilizers. *International Journal of Pharmacy and Pharmaceutical Sciences*, 8(2), 37–40.
- Argaw, A., Mekonnen, E., & Muleta, D. (2015). Agronomic efficiency of N of common bean (Phaseolus vulgaris L.) in some representative soils of Eastern Ethiopia. *Cogent Food and Agriculture*, 1(1). https://doi.org/10.1080/23311932.2015.1074790
- Asadu, C. O., Ezema, C. A., Ekwueme, B. N., Onu, C. E., Onoh, I. M., Adejoh, T., Ezeorba, T. P. C., Ogbonna, C. C., Otuh, P. I., Okoye, J. O., & Emmanuel, U. O. (2024). Enhanced efficiency fertilizers: Overview of production methods, materials used, nutrients release mechanisms, benefits and considerations. *Environmental Pollution and Management*, 1(March), 32–48. https://doi.org/10.1016/j. epm.2024.07.002
- Carneiro, R., Duncan, S., O'Keefe, S., Yu, D., Huang, H., Yin, Y., Neill, C., Zhang, B., Kuhar, T., Rideout, S., Reiter, M., Ross, J., Chen, P., & Gillen, A. (2021). Utilizing consumer perception of edamame to guide new variety development. *Frontiers in Sustainable Food Systems*, 4(January), 1–14. https://doi.org/10.3389/fsufs.2020.556580
- Congreves, K. A., Otchere, O., Ferland, D., Farzadfar, S., Williams, S., & Arcand, M. M. (2021). Nitrogen use efficiency definitions of today and tomorrow. *Frontiers in Plant Science*, *12*(June), 1–10. https://doi.org/10.3389/fpls.2021.637108

- 10. Da Costa, P. B., Granada, C. E., Ambrosini, A., Moreira, F., De Souza, R., Dos Passos, J. F. M., Arruda, L., & Passaglia, L. M. P. (2014). A model to explain plant growth promotion traits: A multivariate analysis of 2,211 bacterial isolates. *PLoS ONE*, 9(12), 1–25. https://doi.org/10.1371/journal. pone.0116020
- De Sousa Lima, J. R., De Goes, M. da C. C., Hammecker, C., Antonino, A. C. D., De Medeiros, É. V., De Sá Barretto Sampaio, E. V., De Barros Silva Leite, M. C., Da Silva, V. P., De Souza, E. S., & Souza, R. (2021). Effects of poultry manure and biochar on acrisol soil properties and yield of common bean. a short-term field experiment. *Agriculture (Switzerland)*, *11*(4), 1–11. https://doi.org/10.3390/agriculture11040290
- Dhaliwal, S. S., Sharma, V., Shukla, A. K., Gupta, R. K., Verma, V., Kaur, M., Behera, S. K., & Singh, P. (2023). Residual effect of organic and inorganic fertilizers on growth, yield and nutrient uptake in wheat under a Basmati Rice–Wheat cropping system in North-Western India. *Agriculture (Switzerland)*, 13(3). https://doi.org/10.3390/agriculture13030556
- Dobermann, A. (2007). Fertilizer Best Management Practices General Principles. In Fertilizer Best Management Practices: General Principles, Strategy for Their Adoption and Voluntary Initiatives Versus Regulations. https://digitalcommons. unl.edu/agronomyfacpub/1442
- 14. Fathi, A. (2022). Role of nitrogen (N) in plant growth, photosynthesis pigments, and N use efficiency: A review. *Agrisost*\, 28(October). https:// doi.org/10.5281/zenodo.7143588
- 15. Fhonna, T. N., Melati, M., & Aziz, S. A. (2023). Twice applications of laying hens manure produced the highest yield of organic okra pods (Abelmoschus esculentus L. Moench.). *Journal of Tropical Crop Science*, *10*(03), 176–185. https://doi.org/10.29244/ jtcs.10.03.176-185
- Gajanand, Kumar, P., & Deelip, B. S. (2020). Nutrient use efficiency: An overview. *Just Agriculture*, *1*(3), 1–4.
- Hawkesford, M., Horst, W., Kichey, T., Lambers, H., Schjoerring, J., Moller, I. S., & White, P. (2012). Functions of macronutrients. In P. Marschner (Ed.), *Mineral Nutrition of Higher Plants.* (3rd ed., p. 135190). Academic Press.
- 18. He, P., Li, S., Jin, J., Wang, H., Li, C., Wang, Y., & Cui, R. (2009). Performance of an optimized nutrient management system for double-cropped wheat-maize rotations in North-Central China. *Agronomy Journal*, 101(6), 1489–1496. https://doi. org/10.2134/agronj2009.0099
- Htwe, A. Z., Moh, S. M., Moe, K., & Yamakawa, T. (2019). Biofertilizer production for agronomic application and evaluation of its symbiotic

effectiveness in soybeans. *Agronomy*, 9(4), 1–16. https://doi.org/10.3390/agronomy9040162

- 20. Jabary, M., Melati, M., & Aziz, S. A. (2023). Penentuan dosis dupuk kandang ayam untuk produksi paria secara organik. Determination of chicken manure dosage for bitter gourd production organically. *Prosiding Seminar Nasional PERHORTI*, 37–46.
- 21. Jansa, J., Oberholzer, H.-R., & Egli, S. (2009). Environmental determinants of the arbuscular mycorrhizal fungal infectivity of Swiss agricultural soils. *European Journal of Soil Biology*, 45(5–6), 400–408. https://doi.org/https://doi.org/10.1016/j. ejsobi.2009.07.004
- 22. Javed, T., Indu, I., Singhal, R. K., Shabbir, R., Shah, A. N., Kumar, P., Jinger, D., Dharmappa, P. M., Shad, M. A., Saha, D., Anuragi, H., Adamski, R., & Siuta, D. (2022). Recent advances in agronomic and physio-molecular approaches for improving nitrogen use efficiency in crop plants. *Frontiers in Plant Science*, 13(April), 1–21. https://doi.org/10.3389/ fpls.2022.877544
- 23. Jeong, H., Choi, S. K., Ryu, C. M., & Park, S. H. (2019). Chronicle of a soil bacterium: Paenibacillus polymyxa E681 as a tiny guardian of plant and human health. *Frontiers in Microbiology*, 10, 1–16. https://doi.org/10.3389/fmicb.2019.00467
- 24. Kobierski, M., Bartkowiak, A., Lemanowicz, J., & Piekarczyk, M. (2017). Impact of poultry manure fertilization on chemical and biochemical properties of soils. *Plant, Soil and Environment*, 63(12), 558–563. https://doi.org/10.17221/668/2017-PSE
- 25. Kumar, S., Diksha, Sindhu, S. S., & Kumar, R. (2022). Biofertilizers: An ecofriendly technology for nutrient recycling and environmental sustainability. *Current Research in Microbial Sciences*, 3(July 2021), 100094. https://doi.org/10.1016/j. crmicr.2021.100094
- 26. Lanna, N. B. L., Silva, P. N. L., Colombari, L. F., Corrêa, C. V., & Cardoso, A. I. I. (2018). Residual effect of organic fertilization on radish production. *Horticultura Brasileira*, 36(1), 47–53. https://doi. org/10.1590/S0102-053620180108
- 27. Lin, Y., van Santen, E., & Watts, D. (2016). The effect of poultry litter application on agricultural production: a meta-analysis of crop yield, nutrient uptake and soil fertility. *Conference* on Applied Statistics in Agriculture. https://doi. org/10.4148/2475-7772.1494
- 28. Lozano-Contreras, M. G., Ramírez-Jaramillo, G., & Ramírez-Silva, J. H. (2021). Interaction effects between organic fertilizers and biofertilizers on the growth of Stevia rebaudiana Bertoni. *OALib*, 08(11), 1–13. https://doi.org/10.4236/oalib.1108065
- 29. Luo, L., Zhang, Y., & Xu, G. (2020). How does nitrogen shape plant architecture? *Journal of Experimental Botany*, *71*(15), 4415–4427. https://doi.

org/10.1093/jxb/eraa187

- Magana, R. E. (2020). Enrichment of Organic Fertilizer with Plant Growth-Promoting Bacteria: The Effect on Yield and Quality of Okra Fruit. Master Thesis. Bogor Agricultural University, Indonesia.
- 31. Magana, R. E., Melati, M., Purnamawati, H., & Pratiwi, E. (2020). Enrichment of organic manure with plant growth promoting rhizobacteria improved the root and shoot growth of okra (Abelmoschus esculentus L. Moench.). *Journal of Tropical Crop Science*, 7(03), 137–147. https://doi.org/10.29244/ jtcs.7.03.137-147
- 32. Malusà, E., Pinzari, F., & Canfora, L. (2016). Efficacy of biofertilizers: challenges to improve crop production. In D. Singh, H. Singh, & R. Prabha (Eds.), *Microbial Inoculants in Sustainable Agricultural Productivity* (pp. 17–40). Springer, New Delhi. https://doi.org/10.1007/978-81-322-2644-4\_2
- 33. Mandic, V., Krnjaja, V., Tomic, Z., Bijelic, Z., Simic, A., Muslic, D. R., & Gogic, M. (2015). Nitrogen fertilizer influence on wheat yield and use efficiency under different environmental conditions. *Chilean Journal of Agricultural Research*, 75(1), 92–97. https://doi.org/10.4067/ S0718-58392015000100013
- 34. Marschner, P., Crowley, D., & Yang, C. H. (2004). Development of specific rhizosphere bacterial communities in relation to plant species, nutrition and soil type. *Plant and Soil*, 261(1–2), 199–208. https:// doi.org/10.1023/B:PLSO.0000035569.80747.c5
- 35. Melati, M., Ari, A. N. H. G., & Aziz, S. A. (2021). Plant growth and morphological characters of Sonchus arvensis L. from different chicken manure rates and harvest intervals with ratooning practices. *Journal of Tropical Crop Science*, 8(02), 60–70. https:// doi.org/10.29244/jtcs.8.02.60-70
- 36. Mishra, N., Singh, A., Kumar, P., & Singh, U. (2023). Fertilizer use efficiency. In K. Baboo, S. T. Zaidi, M. M. Kumar, & P. K. Shukla (Eds.), *Impetus Agriculture* (1st ed., Issue June, pp. 114–122). Weser Books.
- 37. Mitter, E. K., Tosi, M., Obregón, D., Dunfield, K. E., & Germida, J. J. (2021). Rethinking crop nutrition in times of modern microbiology: Innovative biofertilizer technologies. *Frontiers in Sustainable Food Systems*, 5(February), 1–23. https://doi. org/10.3389/fsufs.2021.606815
- 38. Mujeeb, F., Ali, A., Qureshi, M. A., Sarwar, G., Ali, M. A., Javed, H., Rafique, M., Ijaz, F., & Ehsan, S. (2022). Comparative study of application methods of biofertilizer for growth enhancement of cereals. 7, 45–53.
- Nosheen, S., Ajmal, I., & Song, Y. (2021). Microbes as biofertilizers, a potential approach for sustainable crop production. *Sustainability (Switzerland)*,

13(4), 1-20. https://doi.org/10.3390/su13041868

- 40. Ortega Pérez, R., Nieto García, J. C., Gallegos-Cedillo, V. M., Domene Ruiz, M. Á., Santos Hernández, M., Nájera, C., Miralles Mellado, I., & Diánez Martínez, F. (2023). Biofertilizers enriched with PGPB improve soil fertility and the productivity of an intensive tomato crop. *Agronomy*, *13*(9), 1–13. https://doi.org/10.3390/agronomy13092286
- 41. Parnell, J. J., Berka, R., Young, H. A., Sturino, J. M., Kang, Y., Barnhart, D. M., & Dileo, M. V. (2016). From the lab to the farm: An industrial perspective of plant beneficial microorganisms. *Frontiers in Plant Science*, 7(AUG2016), 1–12. https://doi. org/10.3389/fpls.2016.01110
- 42. Rasool, A., Ghani, A., Nawaz, R., Ahmad, S., Shahzad, K., Rebi, A., Ali, B., Zhou, J., Ahmad, M. I., Tahir, M. F., Alwahibi, M. S., Elshikh, M. S., & Ercisli, S. (2023). Effects of poultry manure on the growth, physiology, yield, and yield-related traits of maize varieties. *ACS Omega*, 8(29), 25766–25779. https://doi.org/10.1021/acsomega.3c00880
- 43. Roggatz, U., McDonald, A. J. S., Stadenberg, I., & Schurr, U. (1999). Effects of nitrogen deprivation on cell division and expansion in leaves of Ricinus communis L. *Plant, Cell and Environment*, 22(1), 81–89. https://doi.org/10.1046/j.1365-3040.1999.00383.x
- 44. Sahu, B., Singh, J., Shankar, G., & Pradhan, A. (2018). Pseudomonas fluorescens PGPR bacteria as well as biocontrol agent: A review. *International Journal of Chemical Studies*, 6(2), 1–7.
- 45. Samantaray, A., Chattaraj, S., Mitra, D., Ganguly, A., Kumar, R., Gaur, A., Mohapatra, P. K. D., Santos-Villalobos, S. de los, Rani, A., & Thatoi, H. (2024). Advances in microbial based bio-inoculum for amelioration of soil health and sustainable crop production. *Current Research in Microbial Sciences*, 7(June), 100251. https://doi.org/10.1016/j.crmicr.2024.100251
- 46. Sardans, J., & Peñuelas, J. (2021). Potassium control of plant functions: Ecological and agricultural implications. In *Plants* (Vol. 10, Issue 2). https://doi. org/10.3390/plants10020419
- 47. Schütz, L., Gattinger, A., Meier, M., Müller, A., Boller, T., Mäder, P., & Mathimaran, N. (2018). Improving crop yield and nutrient use efficiency via biofertilization— A global meta-analysis. *Frontiers in Plant Science*, 8(January). https://doi.org/10.3389/fpls.2017.02204
- 48. Sharma, S., Kaur, G., Singh, P., Alamri, S., Kumar, R., & Siddiqui, M. H. (2022). Nitrogen and potassium application effects on productivity, profitability and nutrient use efficiency of irrigated wheat (Triticum aestivum L.). *PLoS ONE*, *17*(5 May), 1–29. https://doi.org/10.1371/journal.pone.0264210
- 49. Singh, S. P., Singh, S., Dubey, A. N., & Rajput, R. K. (2020). Biofertilizers and plant growth regulators as key player in sustainability. In *Three Major Dimensions of Life: Environment, Agriculture and*

*Health* (pp. 12–18).

- 50. Soremi, A., Adetunji, M., Adejuyigbe, C., Bodunde, J., & Azeez, J. (2017). Effects of poultry manure on some soil chemical properties and nutrient bioavailability to soybean. *Journal of Agriculture and Ecol*ogy Research International, 11(3), 1–10. https://doi. org/10.9734/jaeri/2017/32419
- 51. Suntari, Ghulamahdi, M., & Melati, M. (2023). Relay-cropping soybean-maize in saturated soil culture increases efficiency of land use and nitrogen fertilizer. *Indonesian Journal of Agronomy*, 51(1), 91–100. https://doi.org/10.24831/ija.v51i1.44905
- 52. Thilakarathna, M. S., & Raizada, M. N. (2017). A meta-analysis of the effectiveness of diverse rhizobia inoculants on soybean traits under field conditions. *Soil Biology and Biochemistry*, 105, 177–196. https://doi.org/https://doi.org/10.1016/j. soilbio.2016.11.022
- 53. Timmusk, S., Grantcharova, N., & Wagner, E. G. H. (2005). Paenibacillus polymyxa invades plant roots and forms biofilms. *Applied and Environmental Microbiology*, 71(11), 7292–7300. https://doi. org/10.1128/AEM.71.11.7292-7300.2005
- 54. Weber, N. F., Herrmann, I., Hochholdinger, F., Ludewig, U., & Neumann, G. (2018). PGPR-induced growth stimulation and nutrient acquisition in maize: Do root hairs matter? *Scientia Agriculturae Bohemica*, 49(3), 164–172. https://doi.org/10.2478/ sab-2018-0022
- 55. Yang, J., Lan, L., Jin, Y., Yu, N., Wang, D., & Wang, E. (2022). Mechanisms underlying legume-rhizobium symbioses. *Journal of Integrative Plant Biology*, 64(2), 244–267. https://doi.org/10.1111/jipb.13207
- 56. Yu, D., Lin, T., Sutton, K., Lord, N., Carneiro, R., Jin, Q., Zhang, B., Kuhar, T., Rideout, S., Ross, J., Duncan, S., Yin, Y., Wang, H., & Huang, H. (2021). Chemical compositions of edamame genotypes grown in different locations in the US. *Frontiers in Sustainable Food Systems*, 5(February), 1–14. https://doi.org/10.3389/fsufs.2021.620426
- 57. Zeipina, S., Alsina, I., & Lepse, L. (2017). Insight in edamame yield and quality parameters: A review. *Research for Rural Development*, 2, 40–44. https:// doi.org/10.22616/rrd.23.2017.047
- 58. Zhang, Y., Liu, J., Zhang, J., Liu, H., Liu, S., Zhai, L., Wang, H., Lei, Q., Ren, T., & Yin, C. (2015). Row ratios of intercropping maize and soybean can affect agronomic efficiency of the system and subsequent wheat. *PLoS ONE*, *10*(6), 1–16. https://doi. org/10.1371/journal.pone.0129245
- 59. Zhao, Z., Mao, Y., Gao, S., Lu, C., Pan, C., & Li, X. (2023). Organic carbon accumulation and aggregate formation in soils under organic and inorganic fertilizer management practices in a rice–wheat cropping system. *Scientific Reports*, *13*(1), 1–12. https://doi. org/10.1038/s41598-023-30541-y